



ISSN 2231-3478

(Print)

JUSPS-B Vol. 29(10), 295-302 (2017). Periodicity-Monthly

Section B

(Online)



ISSN 2319-8052

9 772319 805003



Estd. 1989

JOURNAL OF ULTRA SCIENTIST OF PHYSICAL SCIENCES

An International Open Free Access Peer Reviewed Research Journal of Physical Sciences

website:- www.ultrascientist.org**Synthesis and physicochemical characteristics of Nickel (II), Copper (II), Zinc (II), Cadmium (II) and Tin (II) complexes of schiff base with 1, 2, 4-triazole**POOJA¹, VINITA GUPTA², SANCHITA SINGH³ and Y.K. GUPTA^{4*}¹Department of Chemistry, Janta Vidya Mandir Ganpat Rai Rasiwasia (J.V.M.G.R.R.) P.G. College, Charkhi Dadri, Haryana (India)^{2,3}Department of Chemistry, Agra College, Agra, U.P., (India)⁴Head Department of Chemistry, B K Birla Institute of Engineering and Technology, Pilani, Rajasthan, (India)Corresponding Author Email:- ykgbkbiet@rediffmail.com, ykgbkbiet123@gmail.com<http://dx.doi.org/10.22147/jusps-B/291007>**Acceptance Date 13th September, 2017, Online Publication Date 2nd October, 2017****Abstract**

The extreme efforts have been developed to design novel compounds to strains of resistant micro-organisms. On-going search for novel and amazing drug delivery systems is primarily impact of the well-established fact that the convectional dosages are not sufficiently effective in conveying the drug compounds to its site of action and this has required the need to search for more efficient drugs. New five metal complexes of Nickel(II), Copper(II), Zinc(II), Cadmium(II) and Tin(II) with Schiff base ligand 4-(Benzylidene-amino)- 5-pyridin-4-yl-4H-[1,2,4] triazole-3-thiol were synthesized in ethyl alcohol as a solvent. The Synthesized compounds were characterized by different physicochemical techniques such as, elemental analysis C.H.N.S, FT-IR, UV-visible spectrophotometry, ¹H NMR, conductivity and magnetic susceptibility measurements melting point, flame atomic absorption. The Schiff base ligand act as bidentate and coordinated to the metal atoms via the azomethine-N and S atom of thiol group. From the spectral studies, square planar geometry was proposed for the copper complex. The other complexes were proposed to be tetrahedral.

Key words : Metal complexes, Schiff base, bidentate, 1,2,4-triazole.

1. Introduction

Researchers interested in synthesizing and studying the structural aspects of metal complexes containing Sulphur and nitrogen donor ligands because of their increasing physiological importance and the

This is an open access article under the CC BY-NC-SA license (<https://creativecommons.org/licenses/by-nc-sa/4.0>)


active role played by certain metal ions coordinated to them¹⁻⁴. The chemistry of 1,2,4-triazole compounds has attracted widespread attention due to their diverse biological activities, especially as antimicrobial agents⁵. Triazole are heterocyclic compound featuring five member ring of two carbon atoms and three nitrogen atoms as part of the aromatic system⁶. There are two types of triazoles, the 1,2,3-triazoles and the 1,2,4-triazoles^{7,8}. Tautomerism form could occur in triazole⁹. Several compounds containing 1,2,4-triazole ring are well known for drug synthesis¹⁰. 1,2,4-Triazole containing amino group is also important for obtaining various Schiff's bases with well-established antimicrobial properties^[11]. The synthesis of high nitrogen containing heterocyclic systems has been attracting increasing interest over the past decade because of their utility in various applications¹². Schiff's bases are the compounds containing azimethine group (HC=N). They are condensation products of ketones or aldehydes with primary amines and were first reported by Hugo Schiff in 1864^[13]. Schiff's bases of aliphatic aldehydes are relatively unstable and are readily polymerized while those of aromatic aldehydes, having an effective conjugation system, are more stable. Schiff's bases derived from triazole were reported to possess antimicrobial, anti-anxiety, anti-depressant, plant growth regulatory activity^[14]. As ligands, Schiff's bases have been extensively studied. The azomethine group (HC=N), supported by other electron donor groups form stable chelates with many metal ions. Schiff's bases have often been used as chelating ligands in the field of coordination chemistry and their metal complexes are of great interest for many years. It is well known that N and S atoms play a key role in the coordination of metals at the active sites of numerous metallobiomolecules^[15]. The attention Schiff base metal complexes have earned was because they have industrial, antifungal, antibacterial, anticancer and herbicidal applications^[16,17,18]. In this work, we report the preparation of a new series of some of such metal complexes.

2. Experimental

All used chemicals were of analytical analar grade. Metal salts used in this study are Copper acetate Cu (CH₃COO)₂, Cadmium acetate dihydrate Cd (CH₃COO)₂.2H₂O, Tin chloride SnCl₂, Zinc acetate dihydrate Zn (CH₃COO)₂.2H₂O and Nickel nitrate hexahydrate Ni (NO₃)₂.6H₂O. The metal analyses were performed by using GBS-933 Flame plus Atomic Absorption Spectrophotometer. The elemental Analysis (C.H.N.S) of compounds was carried out with EM- 1108. The electronic spectra were recorded on a Shimadzu UV-160 Spectrophotometer. FTIR spectra were recorded using FTIR.8300 Shimadzu spectrophotometer as CsI disc. Magnetic susceptibility measurements were determined using Magnetic Susceptibility Balance Johnson Matthey at room temperature. The molar conductance was measured in ethanol as a solvent at room temperature using WTW conduct meter. Melting point apparatus of Gallen Kamp M.F.B-60 was used for the determination of melting points. The spectra of ¹H NMR were recorded on a Bruker Ultra shield 300 MHZ, using deuterated DMSO-d₆ as the solvent and tetramethyl silane, TMS as the internal standard.

2.1 Synthesis of 4-(Benzylidene-amino)-5-pyridin-4-yl-4H-[1,2,4]triazole-3-thiol (L) :

The Schiff base ligand was prepared by the condensation of respective aldehyde & amino compound. A mixture of 4-Amino-5-pyridin-4-yl-4H-[1,2,4] triazole-3-thiol (1g, 0.005mol), benzaldehyde (0.005mol) and 4-5 drops of glacial acetic acid in ethanol medium was refluxed for 3 hrs^[19]. The resulting solution was cooled to room temperature and the precipitated solid, was filtered, washed with cold ethanol and recrystallized from hot ethanol, **Scheme 1**.



4-(Benzylidene-amino)-5-pyridin-4-yl-4H-[1,2,4]triazole-3-thiol
Scheme 1. General step of preparation of the ligand (L)

2.2 Synthesis of metal complexes:

A solution of the suitable metal salt [Copper (II) acetate, Cadmium (II) acetate dihydrate, Tin (II) chloride, Zinc (II) acetate dihydrate, Nickel (II) nitrate hexahydrate], a hot solution of the ligand were mixed in 2:1 (ligand : metal) molar ratios. The resulting mixture was stirred under reflux for 1h whereupon the crystalline colure complexes precipitated. Thereafter, crystalline complexes slowly separated. The complexes were filtered and washed with hot alcohol. The products were dried at room temperature²⁰.


3. Results and Discussion

The physical analytical data, melting point and elemental analysis of ligand (L) and its complexes are given in **Table 1**. The percent's of Cu(II), Cd(II), Sn(II), Zn(II) and Ni(II) in metal chelates were estimated by flame atomic absorption spectroscopy (F.A.A.S) using GBS-933 Flame Plus Atomic Absorption Spectrophotometer. The calculated values were in good agreement with the experimental values and the melting points were sharp, indicating the purity of the prepared compounds.

3.1 FTIR spectra :

The structure of the prepared ligand (L) and the metal (II) complexes were confirmed by infrared spectroscopy. The characteristic bands in the IR spectra of ligand (L) and its metal (II) complexes are reported in **Table 2**. The disappearance of original bands at 3250, 3213 and 1699 cm⁻¹ due to amino (NH₂) and aldehyde (CHO) groups and, in turn, the appearance of a new band at 1656 cm⁻¹ assigned to the azomethine (-C=N) linkage provided a clue to the Condensation of amine with aldehyde resulting in the formation of the desired Schiff base s²¹. The IR spectra of the Schiff base ligand exhibited bands at 2660, 1629, 1656, 1570, 3032 and

682 cm^{-1} respectively assigned to the vibrations (S-H), (C=N) triazole ring, (C=N) azomethine, (C=C), (C-H) aromatic and the last one is for stretching of (C-S) bond^{22,23}. The tautomerism form could occur in triazole. As a result, the band at 1286 cm^{-1} assigned to the (C=S) vibrations could be observed in the spectra of the ligand²⁴. It is reasonable to expect deprotonation of the ligand molecules before complexation; the complete disappearance of the band due to ν (S-H) in the spectra of the complexes unambiguously supports this view. The Schiff base ligand possessed potential donor sites such as the azomethine linkage (-C=N), and the S atom of thiol group, which have a tendency to coordinate with the metal ions. The azomethine group, appearing at 1656 cm^{-1} in the free ligand, is shifted to lower frequency by 3-12 cm^{-1} in the complexes, and a new band appearing at 528–539 cm^{-1} due to ν (M–N) vibrations indicating the involvement of azomethine nitrogen atom in coordination²⁵. However, the band appearing at 682 cm^{-1} due to (C-S) vibration in the spectra of the ligand shifted to a higher frequency by 2–16 cm^{-1} due to increasing of the bond order of carbon sulphate bond, indicating the participation of the S atom in complexation. This coordination was supported by the appearance of a new band at 427–454 cm^{-1} assigned to (M-S)²⁶. All other bands such as C=C, C-H aromatic remain unchanged in the spectra of the ligand and their corresponding metal complexes because they are not participating in the complexation²⁷. Thus, these results establish that the ligand chelates with metal ions through the deprotonated Sulphur of the (thiol) group and the nitrogen of the Azomethine group as bidentate ligand, **Fig. 1**.



Where M = Cu (II), Cd(II), Sn(II), Zn(II) or Ni(II)

Figure 1. Structure of the complexes

3.2 Nuclear Magnetic Resonance :

The ^1H NMR spectra of free ligand was recorded in DMSO-d_6 . The formation of imine group is established by the loss of signal due to the aldehyde group of benzaldehyde at 9.7 in the ^1H NMR spectra²⁸. ^1H NMR spectrum of ligand shows the signals (ppm) at 7.311 (m, 5H, Ar-H), 8.018, 7.788 (d,d, 4H, Ar-H), 2.522 (s, 1H, SH), 8.518 (s, 1H, N=CH).

3.3 Electronic spectra :

The electronic spectra of the Schiff base ligand and all complexes in 10^{-3} M DMSO solution at room temperature and resulting data are listed in **Table 3**. The UV-Visible spectrum of ligand showed three bands at

263, 307 and 331nm. The former two bands are due to the $\pi-\pi^*$ transitions within the aromatic ring and remain almost unchanged in the spectra of metal complexes while the third band is due to the $n-\pi^*$ transitions of azomethine group and also observed in the spectra of metal complexes with a slight shift to lower wavelength, indicating coordination of the ligand to metal center²⁹.

Table 1 Physical data of new ligand (L) and its metal complexes

Com-pound	Colour	M.P. (°C)	Yield (%)	Molar ratio M:L	Mol. Wt.	Elemental analysis theoretical (Experimental) %				Metal percentage	
						C %	H %	N %	S %	Found	Calc.
L	Yellow	216	67	-	281.4	59.78 (59.75)	3.91 (3.80)	24.91 (24.38)	11.38 (10.20)	-	-
CuL ₂	Dark green	242	88.3	1:2	626.2	53.71 (52.85)	3.51 (3.33)	22.38 (21.74)	10.23 (9.65)	10.15	10.05
CdL ₂	Off white	253	58.8	1:2	675.1	49.82 (50.55)	3.26 (2.93)	20.75 (21.06)	9.48 (8.65)	16.66	15.95
SnL ₂	Yellow	227	74.40	1:2	681.3	49.36 (49.18)	3.23 (3.65)	20.56 (19.74)	9.40 (8.68)	17.43	17.48
ZnL ₂	Pale yellow	193	61.60	1:2	628.1	53.55 (53.19)	3.50 (3.64)	22.31 (21.62)	10.20 (10.51)	10.42	10.36
NiL ₂	Green	238	72.78	1:2	621.3	54.13 (54.66)	3.54 (3.01)	22.55 (22.05)	10.31 (10.78)	9.45	9.83

Table 2. Diagnostic FTIR bands of the ligand and complexes (cm⁻¹)

Compound	Triazole ν C=N	Azomethine ν C=N	ν C-S	ν S-H	ν M-N	ν M-S
L	1629	1656	682	2660	-	-
CuL ₂	1610	1653	694	-	528	436
CdL ₂	1612	1650	693	-	530	454
SnL ₂	1618	1642	682	-	531	430
ZnL ₂	1614	1649	696	-	534	427
NiL ₂	1614	1647	698	-	539	451

Table 3. Electronic spectra in DMSO solvent for the prepared ligand and its metal Complexes

Compound	Absorption Bands (nm)	Assignment
L	263, 307, 331	$\pi-\pi^*$, $\pi-\pi^*$, $n-\pi^*$
CuL ₂	263, 307, 326, 494	$\pi-\pi^*$, $\pi-\pi^*$, $n-\pi^*$, $^2T_2 \rightarrow ^2E$
CdL ₂	265, 306, 320	$\pi-\pi^*$, $\pi-\pi^*$, $n-\pi^*$
SnL ₂	263, 307, 329	$\pi-\pi^*$, $\pi-\pi^*$, $n-\pi^*$
ZnL ₂	266, 307, 319	$\pi-\pi^*$, $\pi-\pi^*$, $n-\pi^*$
NiL ₂	263, 307, 323, 591	$\pi-\pi^*$, $\pi-\pi^*$, $n-\pi^*$, $^3T_1(F) \rightarrow ^3T_1(P)$

In the electronic spectra of copper and nickel complexes, one new band appears which may be assigned to d-d transition. The Cu(II) complex exhibits d-d transition band centred at 494 nm, attributable to the $^2T_2 \rightarrow ^2E$

transition suggesting square geometry³⁰. The Ni(II) complex shows a band at 591 nm, which may be assigned to the ${}^3T_1(F) \rightarrow {}^3T_1(P)$ and ${}^3T_1(F) \rightarrow {}^3A_2(F)$ transition, respectively in a tetrahedral environment³¹. Other complexes were diamagnetic as expected for d^{10} ions, so that no (d-d) transition can be expected in the visible region³².

3.3 Magnetic measurements and molar conductivity :

The magnetic moment measurements of compounds were carried out at 25 °C. Magnetic susceptibility measurements provide sufficient data to characterize the structure of the complexes. The magnetic moment for Cu(II) complex was approximately 1.62 B.M. This value refers to square planar structure while the value of Ni(II) complex is approximately 2.74 B.M. leading to suggest a high spin tetrahedral structure³³. Cd(II), Sn(II) and Zn(II) complexes are diamagnetic and there were no magnetic moment recorded in this study. The molar conductivity value of all complexes has been measured in ethanol as a solvent at concentration of 10-3 M at room temperature. This measurement was carried out to determine whether the solution was an electrolyte. **Table 4** shows the molar conductivity measurements of complexes, it shows that all the prepared complexes were found to be nonelectrolyte^[34]. Based on the results obtained from spectral study, magnetic moment and conductivity measurements, all complexes exhibited distorted tetrahedral geometry except Cu(II) complex (distorted square planar)³⁵.

Table 4. Conductivity measurement and magnetic moment of L and its complexes

Compound	Conductivity ($\mu\text{S/cm}$)	Magnetic moment (BM)
L	—	—
CuL ₂	11.4	1.62
CdL ₂	21	0
SnL ₂	14.78	0
ZnL ₂	18.4	0
NiL ₂	22.8	2.74

4. Conclusion

This paper describes the synthesis and characterization of a Schiff base derived from 1,2,4-triazol and its five complexes. The ligand acts as a bidentate and coordinates through azomethine-N, and thiol-S leading to the formation of five member ring chelate. General structures of the complexes are shown in **Fig. 1**. The Cd(II), Sn(II), Zn(II) and Ni(II) complexes probability show tetrahedral geometry, while Cu(II) complex probability shows square planar geometry.

5. Future Scope :

The pharmacological application of transition metal complexes is an under-developed area of research and is full of opportunities for further progress. Basic principles to guide the synthesis and development of transition metal based pharmaceuticals are lacking. Development of new methodologies such as combinatorial chemistry will be helpful in the synthesis of inorganic compounds as therapeutic agents. Similarly, the action of metal complexes in the whole living organisms are expected to differ, in general, from the action of non-metal containing agents, and may offer unique research, diagnostic or therapeutic opportunities.

References

1. Yogesh Kumar Gupta, Vinita Gupta, Sanchita Singh “Synthesis and antimicrobial activities of Co(II), Ni(II)

- and Cu(II) complexes with N-Donor ligand". International Journal of Chemical and Analytical science, (Elsevier) Vol. 04, No. 02, pp 62-66 (2013).
2. S. Jyothi, K. Sreedhar, D. Nagaraju, and S. J. Swamy, "Synthesis and spectral investigation of Co(II), Ni(II), Cu(II) and Zn(II) complexes with novel N4 Ligands" Canadian Chemical Transactions, vol. 3, no. 4, pp. 368–380, (2015).
 3. Vinita Gupta, Sanchita Singh, Y.K.Gupta* "Synthesis, Characterization and antimicrobial activity of Co(II), Ni(II), Cu(II) and Zn(II) complexes N- O-S donor ligands" Asian Journal of Pharmaceutical Analysis, (AJPA) Vol. 04, Issue 04, pp 174-177, (2014).
 4. J. L. Segura, M. J. Mancheño, and F. Zamora, "Covalent organic frameworks based on Schiff-base chemistry: synthesis, properties and potential applications," Chemical Society Review, (2016).
 5. A. K. Ghosh, M. Mitra, A. Fathima et al. "Antibacterial and catecholase activities of Co(III) and Ni(II) Schiff base complexes," Polyhedron, vol. 107, pp. 1–8, (2016).
 6. A. Wahi, A. Singh. Asian J. Biochemical and Pharmaceutical Research, 1, 193-205 (2011).
 7. Y.K. Gupta, S.C. Agarwal, S.P. Madnawat, RamNarain "Synthesis and Antimicrobial Characteristics on Zn(II) and Hg(II) Complexes of 2-Formylpyridine Thiosemicarbazone", International Journal of Research in Chemistry and Environment, Vol. 2, issue 2, pp 153-157, (2012).
 8. D. Bele, I. Singhvi. Asian J. of Biochem and Pharmaceutical Res., 1, 88-100 (2011).
 9. I. Rama and R. Selvameena, "Synthesis, structure analysis, anti-bacterial and in vitro anti-cancer activity of new Schiff base and its copper complex derived from sulfamethoxazole" Journal of Chemical Sciences, vol. 127, no. 4, pp. 671–678 (2015).
 10. Y.K. Gupta, Vinita Gupta, Sanchita Singh "Synthesis, Characterization and Antimicrobial Activity of Nickel and Cobalt Complexes of Schiff bases". International Journal of Chemical and Analytical Science vol.3, No. 8, pp 1537 – 1539, (2012).
 11. A. Singh; O. Pandey; S. Sengupta. Spectrochimica. Acta. Part A, 85, 1-6 (2012).
 12. Y.K. Gupta, S.C. Agarwal, S.P. Madnawat, "Synthesis and Characterization of Schiff Base Complexes of Iron (II) and Nickel (II)", Journal of Ultra Scientist of Physical Sciences, Vol. 24, No 2(B), pp 370-374, (2012).
 13. M. Balouiri, M. Sadiki, and S. K. Ibnsouda "Methods for in vitro evaluating antimicrobial activity: A review," Journal of Pharmaceutical Analysis, vol. 6, pp. 71–79, (2016).
 14. N. Kavitha and P. Anantha Lakshmi, "Synthesis, characterization and thermogravimetric analysis of Co(II), Ni(II), Cu(II) and Zn(II) complexes supported by ONNO tetra dentate Schiff base ligand derived from hydrazino benzoxazine," Journal of Saudi Chemical Society, (2015).
 15. Vinita Gupta, Sanchita Singh, Y.K.Gupta* "Synthesis and antimicrobial activity of Zn (II) and Fe(II) complexes derived from enamine" Research Journal of Engineering and Technology (RJET) Vol. 05, Issue 04, pp 178-183, (2014).
 16. Vinita Gupta, Sanchita Singh, Y.K.Gupta* "Synthesis, characterization and antimicrobial activity of Cu(II), Ni(II) and Co(II) Complexes with N–ODonor Ligand." Asian Journal of Pharmacy and Technology, (AJPTech.) Vol. 04, Issue 04, pp 205-210, (2014).
 17. Y.K. Gupta, S.C. Agarwal, "Physical characterization of some complexes of nitrogen donor compounds", International Journal of Chemical Engineering, Vol. 4, Issue 2, pp 147-158, (2011).
 18. O. E. Sherif and N. S. Abdel-Kader, "DFT calculations, spectroscopic studies, thermal analysis and biological activity of supramolecular Schiff base complexes," Arabian Journal of Chemistry (2015).
 19. N. K. Chaudhary and P. Mishra, "Spectral Investigation and in Vitro Antibacterial Evaluation of NiII and CuII Complexes of Schiff Base Derived from Amoxicillin and 4-Formylthiophene (4 ft)," Journal of Chemistry, vol. 2015, Article ID 136285 (2015).
 20. Y.K. Gupta* Vinita Gupta, Sanchita Singh, "Synthesis, Characterization and Antimicrobial Activity of Cu(II), Co(II), Ni(II) and Zn(II), Complexes with O-N donor ligand ." International Journal of Technology (IJT) Vol. 04, Issue 02, pp 283-286, (2014).

21. M. Hanif; Z. Chohan. *Appl. Organometal. Chem.*, Vol. 27, pp 36-44, (2013).
22. Vinita Gupta, Sanchita Singh, Y.K. Gupta* "Synthesis and Antimicrobial Activities of Zn (II) Complex of 2,5-diamino-1,3,4-thiadiazole". *Research Journal of Sciences and Technology* Vol. 05, Issue 04, pp 462-465, (2013).
23. L. Saghatforoush; A. Aminkhani; S. Ershad; G. Karimnezhad; S. Ghammamy; R. Kabiri. *Molecules*, Vol. 13, pp 804-811, (2008).
24. J. Gowda; A. Khader; B. Kalluraya; P. Shree; A. Shabaraya. *European Journal of Medicinal Chemistry*, Vol. 46, pp 4100-4106, (2011).
25. Y.K. Gupta, S.C. Agarwal, S.P. Madnawat, Ram Narain "Synthesis and Antimicrobial Activities of Mn (II) and Fe(III) with N- S donor Ligand" *Research Journal of Chemistry and Environment*, Vol. 16, No 2, pp 48-51, (2012).
26. R. Haddad; E. Yousif; A. Ahmed. *Springer Plus*, Vol. 02, pp 510-516, (2013).
27. Y.K. Gupta, S.C. Agarwal, S.P. Madnawat, "Synthesis, Characterization and Antimicrobial Studies on Some Bivalent Metal Complexes", *Journal of Chemical and Pharmaceutical Research*, Vol. 4, Issue 1, pp 746-750, (2012).
28. Z. Shaghaghi. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, Vol. 131, pp 67-71, (2014).
29. J. Anaconda; J. Rodriguez; J. Camus. "Spectrochimica Acta Part A": *Molecular and Biomolecular Spectroscopy*, 129, 96-102, (2014).
30. Y.K. Gupta, S.C. Agarwal, "Synthesis, physico-chemical and antimicrobial studies of Cr (III), Mn (III) and Fe (III) Schiff base metal complexes", *International Journal of Chemical and Analytical Science*, Vol. 2, No 11, pp 1256-1259, (2011).
31. Y. Win; E. Yousif; A. Majeed; S. Ha. *Asian J. Chem*, Vol. 23(11), pp 5009-5012, (2011).
32. Y.K. Gupta, S.C. Agarwal, S.P. Madnawat, Ram Narain "Synthesis, Characterization and Antimicrobial Studies of Some Transition Metal Complexes of Schiff Bases", *Research Journal of chemical science*, vol. 2, Issue 4, pp 68-71, (2012).
33. H. Ibraheem; H. Adel; A. Ahmed; N. Salih; J. Salimon; A. Graisa; Y. Farina; E. Yousif. *Journal of Al-Nahrain University*, Vol. 13 (1), pp 43-47, (2010).
34. A. Majeed; *Al Mustansiriya J. Sci.*, Vol. 21(5), pp 195-204, (2010).
35. Y. Foo; E. Yousif; S. Tiong; A. Majeed. *Asian J. Chem.*, Vol. 25(8), pp 4203-4206, (2013).